ERIC H. HOLMI al. Serial No. Not Assigned Page 6

REMARKS

The present amendment cancels claims 1-47 and 60-62 without prejudice to Applicants' right to prosecute which are directed to other patentably distinct groups disclosed and claimed in the present application. With entry of the present amendment claims 48-52 and new claims 63-75 will be pending. The specification has been amended to include information relating to the co-pending parent application, to cancel pages i-iii comprising a Table of Contents, and to incorporate the American Type Culture Collection Accession numbers. Further, claims 48-52 have been amended to remove the multiple dependency. New claims 63-75 have been added to encompass the subject matter of the multiple dependency. No new matter is believed to be added by the above amendments.

CONCLUSION

In view of the foregoing, Applicants believe all claims now pending in this Application are in condition for allowance. The issuance of a formal Notice of Allowance at an early date is respectfully requested. If the Examiner believes a telephone conference would expedite prosecution of this application, please telephone the undersigned at 206-467-9600.

Respectfully submitted,

Dated: / November 2001

Brian W. Poor Reg. No. 32,928

TOWNSEND and TOWNSEND and CREW LLP

Two Embarcadero Center, 8th Floor San Francisco, California 94111-3834

Tel: (206) 467-9600 Fax: (415) 576-0300

BWP:ww SE 5010162 v1 ERIC H. HOLME. al. Serial No. Not Assigned Page 7

APPENDIX

VERSION WITH MARKINGS TO SHOW CHANGES MADE

IN THE SPECIFICATION:

Microorganisms

Accession Number

CAT-RFT-pPROTA in E. coli INVα

[—]<u>207225</u>

FL-RFT-pcDNA3 in E. coli DH5α

[—]<u>207224</u>

IN THE CLAIMS:

48. (Amended) A method for the preparative synthesis of a molecule comprising Fuc $\alpha 1 \rightarrow 2$ Gal $\beta 1 \$ $\rightarrow 3$ GalNAc, said method comprising contacting [the]an isolated or purified [rat] $\alpha 1 \rightarrow 2$ fucosyltransferase [of Claim 1, 2, 3, 4, 5, 6, or 8] comprising an amino acid sequence as depicted in Figure 5 (SEQ ID NO: 8) with GDP-fucose and a molecule having a terminal Gal $\beta 1 \rightarrow 3$ GalNAc moiety and recovering [a]the molecule comprising Fuc $\alpha 1 \rightarrow 2$ Gal $\beta 1 \rightarrow 3$ GalNAc.

2 3

1

2

3

4

5

6

1

glycoprotein, glycolipoprotein or free oligosaccharide comprising Fuc $\alpha 1 \rightarrow 2Gal\beta 1 \rightarrow 3GalNAc$, said method comprising contacting [the]an isolated or purified [rat $\alpha 1 \rightarrow 2fucosyltransferase$ of

(Amended) A method for the preparative synthesis of a glycolipid,

4

claim 1, 2, 3, 4, 5, 6 or 8]protein comprising an amino acid sequence as depicted in Figure 5

5

oligosaccharaide having a terminal Galβ1→3GalNAc moiety and recovering [a]the glycolipid,

(SEQ ID NO: 8) with GDP-fucose and a glycolipid, glycoprotein, glycolipoprotein or free

7

glycoprotein, glycolipoprotein or free oligosaccharaide comprising

8

Fuc $\alpha 1 \rightarrow 2Gal\beta 1 \rightarrow 3GalNAc$.

49.

1 2

3

50. (Amended) The method according to Claim 49 wherein the [rat] α1→2fucosyltransferase is contacted with an oligosaccharide comprising a terminal Galβ1→3GalNAc moiety.

51. (Amended) A method for the preparative synthesis of fucosyl-GM₁ comprising contacting [the]an isolated or purified [rat] $\alpha 1 \rightarrow 2$ fucosyltransferase [of Claim 1, 2, 3, 4, 5, 6 or 8]comprising an amino acid sequence as depicted in Figure 5 (SEQ ID NO: 8) with GDP-fucose and the ganglioside GM₁ and recovering fucosyl-GM₁.

- 52. (Amended) A method for the preparative synthesis of a molecule comprising Fuc $\alpha 1 \rightarrow 2$ Gal $\beta 1 \rightarrow 3$ GalNAc, said method comprising contacting [the isolated or purified rat]a recombinant $\alpha 1 \rightarrow 2$ fucosyltransferase [of Claim 33, 36, 39, or 42] or [the]a cellular fraction [of Claim 34, 37, 40, or 43]of a recombinant cell containing a vector having a nucleotide sequence that encodes and expresses an amino acid sequence as depicted in Figure 5 (SEQ ID NO: 8) and having $\alpha 1 \rightarrow 2$ fucosyltransferase activity, with GDP-fucose and a molecule having a terminal Gal $\beta 1 \rightarrow 3$ GalNAc moiety and recovering a molecule comprising Fuc $\alpha 1 \rightarrow 2$ Gal $\beta 1 \rightarrow 3$ GalNAc.
- 53. (Amended) A method for the preparative synthesis of a glycolipid, glycoprotein, glycolipoprotein or free oligosaccharide comprising Fucα1→2Galβ1→3GalNAc, said method comprising contacting [the]an isolated or purified recombinant produced rat α1→2fucosyltransferase [of Claim 33, 36, 39, or 42] or [the]a cellular fraction [of Claim 34, 37, 40, or 43]of a recombinant cell containing a vector having a nucleotide sequence as depicted as SEQ ID NO: 7 and having α1→2 fucosyltransferase activity, with GDP-fucose and a glycolipid, glycoprotein, glycolipoprotein or oligosaccharide having a terminal Galβ1→3GalNAc moiety and recovering a glycolipid, glycoprotein, glycolipoprotein or free oligosaccharide comprising Fucα1→2Galβ1→3GalNAc.